Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 2000

Contents

Editorials

Nerup J., Halban P. A.: Sponsor, leader and lobbyist: The European Foundation for the Study of Diabetes (EFSD) and the future of diabetes research in Europe 1453
Riis P.: Obligatory information on patient-oriented end point outcome 135
Waldhäusl W. K.: The 20 th century – the frame that we leave 1

Reviews

Ahrén B.: Autonomic regulation of islet hormone secretion – Implications for health and disease 393
Chan N. N., Vallance P., Colhoun H. M.: Nitric oxide and vascular responses in Type I diabetes 137
Flyvbjerg A.: Putative pathophysiological role of growth factors and cytokines in experimental diabetic kidney disease 1205
Frühbeck G., Salvador J.: Relation between leptin and the regulation of glucose metabolism 3
Gebre-Medhin S., Olofsson C., Mulder H.: Islet amyloid polypeptide in the islets of Langerhans: friend or foe? 687
Knip M., Douek I. F., Moore W. P. T., Gillmor H. A., McLean A. E. M., Bingley P. J., Gale EAM for the ENDIT Group: Safety of high-dose nicotinamide: a review 1337
Nonogaki K.: New insights into sympathetic regulation of glucose and fat metabolism 533
Scharfmann R.: Control of early development of the pancreas in rodents and humans: implications of signals from the mesenchyme 1083
Slatter D. A., Bolton C. H., Bailey A. J.: The importance of lipid-derived malondialdehyde in diabetes mellitus 550
Standl E., Schnell O.: A new look at the heart in diabetes mellitus: from ailing to failing 1455
Vinik A. I., Park T. S., Stansberry K. B., Pittenger G. L.: Diabetic neuropathies 957
Wollheim C. B.: Beta-cell mitochondria in the regulation of insulin secretion: a new culprit in Type II diabetes 265
Zierath J. R., Krook A., Wallberg-Henriksson H.: Insulin action and insulin resistance in human skeletal muscle 821

Articles

Accardo-Palumbo A., Triolo G., Colonna-Romano G., Potestio M., Carbone M., Ferrante A., Giardina E., Caimi G., Triolo G.: Glucose-induced loss of glycosyl-phosphatidylinositol-anchored membrane regulators of complement activation (CD59, CD55) by in vitro cultured human umbilical vein endothelial cells 1039
Acerini C. L., Cheetham T. D., Edge J. A., Dunger D. B.: Both insulin sensitivity and insulin clearance in children and young adults with Type I (insulin-dependent) diabetes vary with growth hormone concentrations and with age 61
Aiston S., Peak M., Agius L.: Impaired glycogen synthesis in hepatocytes from Zucker fatty *fa/fa* rats: the role of increased phosphorylase activity 589
Akimoto Y., Kreppel L. K., Hirano H., Hart G. W.: Increased 0-GlcNAc transferase in pancreas of rats with streptozotocin-induced diabetes 1239
Albareda M., Rodríguez-Espinosa J., Murugo M., de Leiva A., Corcoy R.: Assessment of insulin sensitivity and beta-cell function from measurements in the fasting state and during an oral glucose tolerance test 1507

Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 2000

Contents

Editorials

Nerup J., Halban P. A.: Sponsor, leader and lobbyist: The European Foundation for the Study of Diabetes (EFSD) and the future of diabetes research in Europe 1453
Riis P.: Obligatory information on patient-oriented end point outcome 135
Waldhäusl W. K.: The 20 th century – the frame that we leave 1

Reviews

Ahrén B.: Autonomic regulation of islet hormone secretion – Implications for health and disease 393
Chan N. N., Vallance P., Colhoun H. M.: Nitric oxide and vascular responses in Type I diabetes 137
Flyvbjerg A.: Putative pathophysiological role of growth factors and cytokines in experimental diabetic kidney disease 1205
Frühbeck G., Salvador J.: Relation between leptin and the regulation of glucose metabolism 3
Gebre-Medhin S., Olofsson C., Mulder H.: Islet amyloid polypeptide in the islets of Langerhans: friend or foe? 687
Knip M., Douek I. F., Moore W. P. T., Gillmor H. A., McLean A. E. M., Bingley P. J., Gale EAM for the ENDIT Group: Safety of high-dose nicotinamide: a review 1337
Nonogaki K.: New insights into sympathetic regulation of glucose and fat metabolism 533
Scharfmann R.: Control of early development of the pancreas in rodents and humans: implications of signals from the mesenchyme 1083
Slatter D. A., Bolton C. H., Bailey A. J.: The importance of lipid-derived malondialdehyde in diabetes mellitus 550
Standl E., Schnell O.: A new look at the heart in diabetes mellitus: from ailing to failing 1455
Vinik A. I., Park T. S., Stansberry K. B., Pittenger G. L.: Diabetic neuropathies 957
Wollheim C. B.: Beta-cell mitochondria in the regulation of insulin secretion: a new culprit in Type II diabetes 265
Zierath J. R., Krook A., Wallberg-Henriksson H.: Insulin action and insulin resistance in human skeletal muscle 821

Articles

Accardo-Palumbo A., Triolo G., Colonna-Romano G., Potestio M., Carbone M., Ferrante A., Giardina E., Caimi G., Triolo G.: Glucose-induced loss of glycosyl-phosphatidylinositol-anchored membrane regulators of complement activation (CD59, CD55) by in vitro cultured human umbilical vein endothelial cells 1039
Acerini C. L., Cheetham T. D., Edge J. A., Dunger D. B.: Both insulin sensitivity and insulin clearance in children and young adults with Type I (insulin-dependent) diabetes vary with growth hormone concentrations and with age 61
Aiston S., Peak M., Agius L.: Impaired glycogen synthesis in hepatocytes from Zucker fatty *fa/fa* rats: the role of increased phosphorylase activity 589
Akimoto Y., Kreppel L. K., Hirano H., Hart G. W.: Increased 0-GlcNAc transferase in pancreas of rats with streptozotocin-induced diabetes 1239
Albareda M., Rodríguez-Espinosa J., Murugo M., de Leiva A., Corcoy R.: Assessment of insulin sensitivity and beta-cell function from measurements in the fasting state and during an oral glucose tolerance test 1507

Andersen NA, Larsen C.M., Mandrup-Poulsen T.: TNFα and IFNγ potentiate IL-1β induced mitogen activated protein kinase activity in rat pancreatic islets of Langerhans 1389
Andreelli F., Laville M., Vega N., Riou J.-P., Vidal H.: Regulation of gene expression during severe caloric restriction: lack of induction of p85alpha phosphatidylinositol 3-kinase mRNA in skeletal muscle of patients with Type II (non-insulin-dependent) diabetes mellitus 356
Araki S., Makita Y., Canani L., Ng D., Warram J.H., Krolewski A.S.: Polymorphisms of human paraoxonase 1 gene *(PON1)* and susceptibility to diabetic nephropathy in Type I diabetes mellitus *(Short communications)* 1540
Babawale M.O., Lovat S., Mayhew T.M., Lammiman M.J., James D.K., Leach L.: Effects of gestational diabetes on junctional adhesion molecules in human term placental vasculature 1185
Bacci S., De Cosmo S. , Garruba M., Placentino G., Liuzzi A., Barbano F., Di Giorgio A., Trischitta V., Viberti G.C.: Role of insulin-like growth factor (IGF)-1 in the modulation of renal haemodynamics in Type I diabetic patients 922
Baynes K.C.R., Beeton C.A., Panayotou G., Stein R., Soos M., Hansen T., Simpson H., O'Rahilly S., Shepherd P.R., Whitehead J.P.: Natural variants of human p85α phosphoinositide 3-kinase in severe insulin resistance: a novel variant with impaired insulin-stimulated lipid kinase activity 321
Bernard S., Touzet S., Personne I., Lapras V., Bondon P.J., Berthezène F., Moulin P.: Association between microsomal triglyceride transfer protein gene polymorphism and the biological features of liver steatosis in patients with Type II diabetes 995
Biesenbach G., Margreiter R., Königsrainer A., Bösmüller C., Janko O., Brücke P., Gross C., Zazgornik J.: Comparison of progression of macrovascular diseases after kidney or pancreas and kidney transplantation in diabetic patients with end-stage renal disease 231
Birrell A.M., Heffernan S.J., Ansselin A.D., McLennan S., Church D.K., Gillin A.G., Yue D.K.: Functional and structural abnormalities in the nerves of Type I diabetic baboons: aminoguanidine treatment does not improve nerve function 110
Bonora E., Kiechl S., Oberhollenzer F., Egger G., Bonadonna R.C., Muggeo M., Willeit J.: Impaired glucose tolerance, Type II diabetes mellitus and carotid atherosclerosis: prospective results from the Bruneck Study 156
Boquist S., Hamsten A., Karpe F., Ruotolo G.: Insulin and non-esterified fatty acid relations to alimentary lipaemia and plasma concentrations of postprandial triglyceride-rich lipoproteins in healthy middle-aged men 185
Brabant G., Horn R., von zur Mühlen A., Mayr B., Wurster U., Heidenreich F., Schnabel D., Grüters-Kieslich A., Zimmermann-Belsing T., Feldt-Rasmussen U.: Free and protein bound leptin are distinct and independently controlled factors in energy regulation 438
Branchtein L., Schmidt M.I., Matos M.C.G., Yamashita T., Pousada J.M.D.C., Duncan BB for the Brazilian Gestational Diabetes Study Group: Short stature and gestational diabetes in Brazil 848
Byberg L., McKeigue P.M., Zethelius B., Lithell H.O.: Birth weight and the insulin resistance syndrome: association of low birth weight with truncal obesity and raised plasminogen activator inhibitor-1 but not with abdominal obesity or plasma lipid disturbances 54
Caluwaerts S., Pijnenborg R., Luyten C., Van Assche F.A.: Growth characteristics of diabetic rat ectoplacental cones in vivo and in vitro 939
Carraro M., Mancini W., Artero M., Zennaro C., Faccini L., Candido R., Armini L., Calci M., Carretta R., Fabris B.: Albumin permeability in isolated glomeruli in incipient experimental diabetes mellitus 235
Carvalho E., Eliasson B., Wesslau C., Smith U.: Impaired phosphorylation and insulin-stimulated translocation to the plasma membrane of protein kinase B/Akt in adipocytes from Type II diabetic subjects 1107
Casis O., Gallego M., Iriarte M., Sánchez-Chapula J.A.: Effects of diabetic cardiomyopathy on regional electro-physiologic characteristics of rat ventricle 101
Cassell P.G., Saker P.J., Huxtable S.J., Kousta E., Jackson A.E., Hattersley A.T., Frayling T.M., Walker M., Kopelman P.G., Ramachandran A., Snehelatha C., Hitman G.A., McCarthy M.I.: Evidence that single nucleotide polymorphism in the uncoupling protein 3 *(UCP3)* gene influences fat distribution in women of European and Asian origin 1558
Chowienczyk P.J., Brett S.E., Gopaul N.K., Meeking D., Marchetti M., Russell-Jones D.L., Änggård E.E., Ritter J.M.: Oral treatment with an antioxidant (raxofelast) reduces oxidative stress and improves endothelial function in men with Type II diabetes 974
Collison M., Campbell I.W., Salt I.P., Dominiczak A.F., Connell J.M.C., Lyall H., Gould G.W.: Sex hormones induce insulin resistance in 3T3-L1 adipocytes by reducing cellular content of IRS proteins 1374
Colman P.G., Steele C., Couper J.J., Beresford S.J., Powell T., Kewming K., Pollard A., Gellert S., Tait B., Honeyman M., Harrison L.C.: Islet autoimmunity in infants with a Type I diabetic relative is common but is frequently restricted to one autoantibody 203
Cooper M.E., Thallas V., Forbes J., Scalbert E., Sastra S., Darby I., Soulis T.: The cross-link breaker, *N*-phenacylthiazolium bromide prevents vascular advanced glycation end-product accumulation 660
Crowther N.J., Trusler J., Cameron N., Toman M., Gray I.P.: Relation between weight gain and beta-cell secretory activity and non-esterified fatty acid production in 7-year-old African children: results from the Birth to Ten study 978
De Juan J.A., Moya F.J., Ripodas A., Bernal R., Fernandez-Cruz A., Fernandez-Durango R.: Changes in the density and localisation of endothelin receptors in the early stages of rat diabetic retinopathy and the effect of insulin treatment 773
De Vriese A.S., Van de Voorde J., Blom H.J., Vanhoutte P.M., Verbeke M., Lameire N.H.: The impaired renal vasodilator response attributed to endothelium-derived hyperpolarizing factor in streptozotocin – induced diabetic rats is restored by 5-methyltetrahydrofolate 1116
Decallonne B., Overbergh L., Casteels K.M., Gysemans C., Bouillon R., Mathieu C.: Streptococcal wall component OK432 restores sensitivity of non-obese diabetic (NOD) thymocytes to apoptotic signals 1302
Delplanque J., Barat-Houari M., Dina C., Gallina P., Clément K., Guy-Grand B., Vasseur F., Boutin P., Froguel P.: Linkage and association studies between the proopiomelanocortin (*POMC*) gene and obesity in caucasian families *(Short communications)* 1554
Devedjian J.C., Pujol A., Cayla C., George M., Casellas A., Paris H., Bosch F.: Transgenic mice overexpressing α_{2A}-adrenoceptors in pancreatic beta-cells show altered regulation of glucose homeostasis 899
Dubois M., Pattou F., Kerr-Conte J., Gmyr V., Vandewalle B., Desreumaux P., Auwerx J., Schoonjans K., Lefebvre J.: Expression of peroxisome proliferator-activated receptor γ (PPAR γ) in normal human pancreatic islet cells 1165
Duvillard L., Pont F., Florentin E., Gambert P., Vergès B.: Significant improvement of apolipoprotein B-containing

lipoprotein metabolism by insulin treatment in patients with non-insulin-dependent diabetes mellitus 27
Edge A. S. B., Spiro R. G.: A specific structural alteration in the heparan sulphate of human glomerular basement membrane in diabetes 1056
Ekbom P., Damm P., Nørgaard K., Clausen P., Feldt-Rasmussen U., Feldt-Rasmussen B., Nielsen L. H., Mølsted-Pedersen L., Mathiesen E. R.: Urinary albumin excretion and 24-hour blood pressure as predictors of pre-eclampsia in Type I diabetes 927
Ellard S., Beards F., Allen L. I. S., Shepherd M., Ballantyne E., Harvey R., Hattersley A. T.: A high prevalence of glucokinase mutations in gestational diabetic subjects selected by clinical criteria 250
Ellerman K. E., Like A. A.: Susceptibility to diabetes is widely distributed in normal class llu haplotype rats 890
Elsner M., Guldbakke B., Tiedge M., Munday R., Lenzen S.: Relative importance of transport and alkylation for pancreatic beta-cell toxicity of streptozotocin 1528
Fewtrell M. S., Doherty C., Cole T. J., Stafford M., Hales C. N., Lucas A.: Effects of size at birth, gestational age and early growth in preterm infants on glucose and insulin concentrations at 9–12 years 714
Flavell D. M., Pineda Torra I., Jamshidi Y., Evans D., Diamond J. R., Elkeles R. S., Bujac S. R., Miller G., Talmud P. J., Staels B., Humphries S. E.: Variation in the PPARα gene is associated with altered function in vitro and plasma lipid concentrations in Type II diabetic subjects 673
Fritsche A., Stefan N., Hardt E., Häring H., Stumvoll M.: Characterisation of beta-cell dysfunction of impaired glucose tolerance: Evidence for impairment of incretin-induced insulin secretion 852
Gambelunghe G., Ghaderi M., Cosentino A., Falorni AD, Brunetti P., Falorni AL, Sanjeevi C. B.: Association of MHC Class I chain-related A *(MIC-A)* gene polymorphism with Type I diabetes 507
Gerritsen J., Dekker J. M., TenVoorde B. J., Bertelsmann F. W., Kostense P. J., Stehouwer C. D. A., Heine R. J., Nijpels G., Heethaar R. M., Bouter L. M.: Glucose tolerance and other determinants of cardiovascular autonomic function: the Hoorn Study 561
Gervaise N., Garrigue M. A., Lasfargues G., Lecomte P.: Triglycerides, apo C3 and Lp B:C3 and cardiovascular risk in Type II diabetes 703
Gilbert R. E., Kelly D. J., Cox A. J., Wilkinson-Berka J. L., Rumble J. R., Osicka T., Panagiotopoulos S., Lee V., Hendrich E. C., Jerums G., Cooper M. E.: Angiotensin converting enzyme inhibition reduces retinal overexpression of vascular endothelial growth factor and hyperpermeability in experimental diabetes 1360
Gottschling-Zeller H., Röhrig K., Hauner H.: Troglitazone reduces plasminogen activator inhibitor-1 expression and secretion in cultured human adipocytes 377
Hadjivassiliou V., Green M. H. L., Green I. C.: Immunomagnetic purification of beta cells from rat islets of Langerhans 1170
Hahn T., Hahn D., Blaschitz A., Korgun E. T., Desoye G., Dohr G.: Hyperglycaemia-induced subcellular redistribution of GLUT1 glucose transporters in cultured human term placental trophoblast cells 173
Hara M., Wang X., Paz V. P., Cox N. J., Iwasaki N., Ogata M., Iwamoto Y., Bell G. I.: No diabetes-associated mutations in the coding region of the hepatocyte nuclear factor-4γ gene *(HNF4G)* in Japanese patients with MODY 1064
Hedrick C. C., Thorpe S. R., Fu M.-X., Harper C. M., Yoo J., Kim S.-M., Wong H., Peters A. L.: Glycation impairs high-density lipoprotein function 312
Heilbronn L. K., Kind K. L., Pancewicz E., Morris A. M., Noakes M., Clifton P. M.: Association of *–3826 G* Variant in uncoupling protein-1 with increased BMI in overweight Australian women 242
Heitzer T., Krohn K., Albers S., Meinertz T.: Tetrahydrobiopterin improves endothelium-dependent vasodilation by increasing nitric oxide activity in patients with Type II diabetes mellitus 1435
Hiilesmaa V., Suhonen L., Teramo K.: Glycaemic control is associated with pre-eclampsia but not with pregnancy-induced hypertension in women it Type I diabetes mellitus 1534
Hoffstedt J., Eriksson P., Hellström L., Rössner S., Rydén M., Arner P.: Excessive fat accumulation is associated with the TNFα –308 G/A promoter polymorphism in women but not in men 117
Holstein P., Ellitsgaard N., Bornefeldt Olsen B., Ellitsgaard V.: Decreasing incidence of major amputations in people with diabetes 844
Hummel M., Bonifacio E., Stern M., Dittler J., Schimmel A., Ziegler A. G.: Development of celiac disease-associated antibodies in offspring of parents with Type I diabetes 1005
Huypens P., Ling Z., Pipeleers D., Schuit F.: Glucagon receptors on human islet cells contribute to glucose competence of insulin release 1012
Hytinantti T. K., Koistinen H. A., Teramo K., Karonen S.-L., Koivisto V. A., Andersson S.: Increased fetal leptin in Type I diabetes mellitus pregnancies complicated by chronic hypoxia 709
Joseph J. W., Kalitsky J., St-Pierre S., Brubaker P. L.: Oral delivery of glucagon-like peptide-1 in a modified polymer preparation normalizes basal glycaemia in diabetic *db/db* mice 1319
Juhl C. B., Schmitz O., Pincus S., Holst J. J., Veldhuis J., Pørksen N.: Short-term treatment with GLP-1 increases pulsatile insulin secretion in Type II diabetes with no effect on orderliness 583
Kahn S. E., Andrikopoulos S., Verchere C. B., Wang F., Hull R. L., Vidal J.: Oophorectomy promotes islet amyloid formation in a transgenic mouse model of Type II diabetes 1309
Kamal A., Biessels G. J., Duis S. E. J., Gispen W. H.: Learning and hippocampal synaptic plasticity in streptozotocin-diabetic rats: interaction of diabetes and ageing 500
Karamanos B., Porta M., Songini M., Metelko Z., Kerenyi Z., Tamas G., Rottiers R., Stevens L. K., Fuller J. H., and the EURODIAB Type I Complications Study Group: Different risk factors of microangiopathy in patients with Type I diabetes mellitus of short versus long duration. The EURODIAB IDDM Complications Study 348
Karlsson M. G. E., Sederholm Lawesson S., Ludvigsson J.: Th1-like dominance in high-risk first-degree relatives of Type I diabetic patients 742
Kelly M. A., Alvi N. S., Croft N. J., Mijovic C. H., Bottazzo G. F., Barnett A. H.: Genetic and immunological characteristics of Type I diabetes mellitus in an Indo-Aryan population 450
Kessler A., Tomas E., Immler D., Meyer H. E., Zorzano A., Eckel J.: Rab11 is associated with GLUT4-containing vesicles and redistributes in response to insulin 1518
Kluge R., Giesen K., Bahrenberg G., Plum L., Ortlepp J. R., Joost H.-G.: Quantitative trait loci for obesity and insulin resistance *(Nob1, Nob2)* and their interaction with the leptin receptor allele *($Lepr^{A720T/T1044I}$)* in New Zealand obese mice 1565
Koch M., Rett K., Volk A., Maerker E., Haist K., Weisser M., Rettig A., Renn W., Häring H. U.: The tumour necrosis factor alpha –238 G → A and –308 G → A promoter polymorphisms are not associated with insulin sensitivity and insulin secretion in young healthy relatives of Type II Diabetic patients 181
Kohara N., Kimura J., Kaji R., Goto Y., Ishii J., Takiguchi M., Nakai M.:

F-wave latency serves as the most reproducible measure in nerve conduction studies of diabetic polyneuropathy: multicentre analysis in healthy subjects and patients with diabetic polyneuropathy 915

Korbutt G. S., Suarez-Pinzon W. L., Power R. F., Rajotte R. V., Rabinovitch A.: Testicular Sertoli cells exert both protective and destructive effects on syngeneic islet grafts in non-obese diabetic mice 474

Kulmala P., Rahko J., Savola K., Vähäsalo P., Veijola R., Sjöroos M., Reunanen A., Ilonen J., Knip M.: Stability of autoantibodies and their relation to genetic and metabolic markers of Type I diabetes in initially unaffected schoolchildren 457

Kusunoki M., Hara T., Tsutsumi K., Nakamura T., Miyata T., Sakakibara F., Sakamoto S., Ogawa H., Nakaya Y., Storlien L. H.: The lipoprotein lipase activator, NO-1886, suppresses fat accumulation and insulin resistance in rats fed a high-fat diet 875

Larsson H., Ahrén B.: Glucose intolerance is predicted by low insulin secretion and high glucagon secretion: outcome of a prospective study in postmenopausal Caucasian women 194

Larsson H., Lindgärde F., Berglund G., Ahrén B.: Prediction of diabetes using ADA or WHO criteria in post-menopausal women: a 10-year follow-up study 1224

Lebrun P., Arkhammar P., Antoine M.-H., Nguyen Q.-A., Bondo Hansen J., Pirotte B.: A potent diazoxide analogue activating ATP-sensitive K^+ channels and inhibiting insulin release 723

Lehto S., Rönnemaa T., Pyörälä K., Laakso M.: Cardiovascular risk factors clustering with endogenous hyperinsulinaemia predict death from coronary heart disease in patients with Type II diabetes 148

Lehtovirta M., Kaprio J., Forsblom C., Eriksson J., Tuomilehto J., Groop L.: Insulin sensitivity and insulin secretion in monozygotic and dizygotic twins 285

Linn T., Santosa B., Grönemeyer D., Aygen S., Scholz N., Busch M., Bretzel R. G.: Effect of long-term dietary protein intake on glucose metabolism in humans 1257

Loviscach M., Rehman N., Carter L., Mudaliar S., Mohadeen P., Ciaraldi T. P., Veerkamp J. H., Henry R. R.: Distribution of peroxisome proliferator-activated receptors (PPARs) in human skeletal muscle and adipose tissue: relation to insulin action 304

Löwel H., Koenig W., Engel S., Hörmann A., Keil U.: The impact of diabetes mellitus on survival after myocardial infarction: can it be modified by drug treatment? 218

Lu X., Yang X.-Y., Howard R. L., Walsh J. P.: Fatty acids modulate protein kinase C activation in porcine vascular smooth muscle cells independently of their effect on de novo diacylglycerol synthesis 1136

Luppi P., Zanone M. M., Hyoty H., Rudert W. A., Haluszczak C., Alexander A. M., Bertera S., Becker D., Trucco M.: Restricted TCR Vβ gene expression and enterovirus infection in Type I diabetes: a pilot study 1484

Maier V. H., Gould G. W.: Long-term insulin treatment of 3T3-L1 adipocytes results in mis-targeting of GLUT4: implications for insulin-stimulated glucose transport 1273

Maran A., Crepaldi C., Trupiani S., Lucca T., Jori E., Macdonald I. A., Tiengo A., Avogaro A., Del Prato S.: Brain function rescue effect of lactate following hypoglycaemia is not an adaptation process in both normal and Type I diabetic subjects 733

Marçais C., Bernard S., Merlin M., Ulhmann M., Mestre B., Rochet-Mingret L., Revol A., Berthezene F., Moulin P.: Severe hypertriglyceridaemia in Type II diabetes: involvement of *apoC-III Sst-I* polymorphism, *LPL* mutations and *apo E3* deficiency 1346

Marfella R., Nappo F., De Angelis L., Siniscalchi M., Rossi F., Giugliano D.: The effect of acute hyperglycaemia on QTc duration in healthy man 571

McEneny J., O'Kane M. J., Moles K. W., McMaster C., McMaster D., Mercer C., Trimble E. R., Young I. S.: Very low density lipoprotein subfractions in Type II diabetes mellitus: alterations in composition and susceptibility to oxidation 485

McGill M., Molyneaux L., Bolton T., Ioannou K., Uren R., Yue D. K.: Response of Charcot's arthropathy to contact casting: assessment by quantitative techniques 481

McLennan S. V., Martell S. Y. K., Yue D. K.: High glucose concentration inhibits the expression of membrane type metalloproteinase by mesangial cells: possible role in mesangium accumulation 642

Meirhaeghe A., Amouyel P., Helbecque N., Cottel D., Otabe S., Froguel P., Vasseur F.: An uncoupling protein 3 gene polymorphism associated with a lower risk of developing Type II diabetes and with atherogenic lipid profile in a French cohort 1424

Mire-Sluis A. R., Gaines Das R., Lernmark Å and Participants of the study: The World Health Organization International Collaborative Study for Islet Cell Antibodies 1282

Mohan V., Ravikumar R., Shanthi Rani S., Deepa R.: Intimal medial thickness of the carotid artery in South Indian diabetic and non-diabetic subjects: the Chennai Urban Population Study (CUPS) 494

Molinete M., Lilla V., Jain R., Joyce P. B. M., Gorr S.-U., Ravazzola M., Halban P. A.: Trafficking of non-regulated secretory proteins in insulin secreting (INS-1) cells 1157

Moriscot C., Pattou F., Kerr-Conte J., Richard M. J., Lemarchand P., Benhamou P. Y.: Contribution of adenoviral-mediated superoxide dismutase gene transfer to the reduction in nitric oxide-induced cytotoxicity on human islets and INS-1 insulin-secreting cells 625

Moustakas A. K., Routsias J., Papadopoulos G. K.: Modelling of the MHC II allele I-A^{g7} of NOD mouse: pH-dependent changes in specificity at pockets 9 and 6 explain several of the unique properties of this molecule 609

Natali A., Toschi E., Camastra S., Gastaldelli A., Groop L., Ferrannini E., on behalf of the European Group for the Study of Insulin Resistance (EGIR): Determinants of postabsorptive endogenous glucose output in non-diabetic subjects 1266

Natali A., Vichi S., Landi P., Severi S., L'Abbate A., Ferrannini E.: Coronary atherosclerosis in Type II diabetes: angiographic findings and clinical outcome 632

New J. P., Hollis S., Campbell F., McDowell D., Burns E., Dornan T. L., Young R. J.: Measuring clinical performance and outcomes from diabetes information systems: an observational study 836

Nolsøe R. L., Kristiansen O. P., Sangthongpitag K., Larsen Z. M., Johannesen J., Karlsen A. E., Pociot F., Nerup J., Verge C. F., Mandrup-Poulsen T.: Complete molecular scanning of the human Fas gene: mutational analysis and linkage studies in families with Type I diabetes mellitus 800

Obrosova I. G., Fathallah L.: Evaluation of an aldose reductase inhibitor on lens metabolism, ATPases and antioxidative defense in streptozotocin-diabetic rats: an intervention study 1048

Ohly P., Dohle C., Abel J., Seissler J., Gleichmann H.: Zinc sulphate induces metallothionein in pancreatic islets of mice and protects against diabetes induced by multiple low doses of streptozotocin 1020

Okamoto S., Kimura K., Kitamura T., Cañas X., Yoshida T., Saito M.: Proinsulin C peptide obviates sympathetically mediated suppression of splenic lymphocyte activity in rats 1512

Olsen H., Länne T.: Reduced capillary hydraulic conductivity in skeletal muscle and skin in Type I diabetes: a possible cause for reduced transcapillary fluid absorption during hypovolaemia 1178

Olsson J., Lindberg G., Gottsäter M., Lindwall K., Sjöstrand Å., Tisell A., Melander A.: Increased mortality in Type II diabetic patients using sulphonylurea and metformin in combination: a population-based observational study 558

Ortsäter H., Liss P., Lund P.E., Åkerman K.E.O., Bergsten P.: Oscillations in oxygen tension and insulin release of individual pancreatic *ob/ob* mouse islets 1313

Osmond D.T.D., Nolan C.J., King R.G., Brennecke S.P., Gude N.M.: Effects of gestational diabetes on human placental glucose uptake, transfer, and utilisation 576

Otabe S., Clement K., Dina C., Pelloux V., Guy-Grand B., Froguel P., Vasseur F.: A genetic variation in the 5′flanking region of the *UCP3* gene is associated with body mass index in humans in interaction with physical activity 245

Otonkoski T., Roivainen M., Vaarala O., Dinesen B., Leipälä J.A., Hovi T., Knip M.: Neonatal Type I diabetes associated with maternal echovirus 6 infection: a case report 1235

Ou D., Mitchell L.A., Metzger D.L., Gillam S., Tingle A.J.: Cross-reactive rubella virus and glutamic acid decarboxylase (65 and 67) protein determinants recognised by T cells of patients with Type I diabetes mellitus 750

Palmer A.J., Weiss C., Sendi P.P., Neeser K., Brandt A., Singh G., Wenzel H., Spinas G.A.: The cost-effectiveness of different management strategies for Type I diabetes: a Swiss perspective 13

Park C.W., Kim J.H., Lee J.W., Kim Y.S., Ahn H.J., Shin Y.S., Kim S.Y., Choi E.J.,Chang Y.S., Bang B.K.: High glucose-induced intercellular adhesion molecule-1 (ICAM-1) expression through an osmotic effect in rat mesangial cells is PKC-NF-κB-dependent 1544

Park Y.S., Kawasaki E., Kelemen K., Yu L., Schiller M.R., Rewers M., Mizuta M., Eisenbarth G.S., Hutton J.C.: Humoral autoreactivity to an alternatively spliced variant of ICA512/IA-2 in Type I diabetes 1293

Pascot A., Després J.P., Lemieux I., Bergeron J., Nadeau A., Prud'homme D., Tremblay A., Lemieux S.: Contribution of visceral obesity to the deterioration of the metabolic risk profile in men with impaired glucose tolerance 1126

Perreault M., Dombrowski L., Marette A.: Mechanism of impaired nitric oxide synthase activity in skeletal muscle of streptozotocin-induced diabetic rats 427

Picard F., Richard D., Timofeeva E., Deshaies Y.: Abnormal insulin and β-adrenergic modulation of lipoprotein lipase during refeeding after prolonged fasting in the Zucker rat 866

Pozzilli P., Pitocco D., Visalli N., Cavallo M.G., Buzzetti R., Crinò A., Spera S., Suraci C., Multari G., Cervoni M., Manca Bitti M.L., Matteoli M.C., Marietti G., Ferrazzoli F., Cassone Faldetta M.R., Giordano C., Sbriglia M., Sarugeri E., Ghirlanda G and the IMDIAB Group: No effect of oral insulin on residual beta-cell function in recent-onset Type I diabetes (the IMDIAB VII) 1000

Price J.A., Fossey S.C., Sale M.M., Brewer C.S., Freedman B.I., Wuerth J.-P., Bowden D.W.: Analysis of the *HNF4α* gene in Caucasian Type II diabetic nephropathic patients 364

Pundziūtė-Lyckå A., Urbonaitė B., Dahlquist G.: Infections and risk of Type I (insulin-dependent) diabetes mellitus in Lithuanian children 1229

Qiao Q., Nakagami T., Tuomilehto J., Borch-Johnsen K., Balkau B., Iwamoto Y., Tajima N., The DECODA Study Group on behalf of the International Diabetes Epidemiology Group: Comparison of the fasting and the 2-h glucose criteria for diabetes in different Asian cohorts 1470

Raskin P., Rappaport E.B., Cole S.T., Yan Y., Patwardhan R., Freed M.I.: Rosiglitazone short-term monotherapy lowers fasting and post-prandial glucose in patients with Type II diabetes 278

Rasmussen S.K., Hansen L., Frevert E.U., Cohen P.T.W., Kahn B.B., Pedersen O.: Adenovirus-mediated expression of a naturally occurring Asp905Tyr variant of the glycogen-associated regulatory subunit of protein phosphatase-1 in L6 myotubes 718

Riemens S.C., Sluiter W.J., Dullaart R.P.F.: Enhanced escape of non-esterified fatty acids from tissue uptake: its role in impaired insulin-induced lowering of total rate of appearance in obesity and Type II diabetes mellitus 416

Rochlitz H., Voigt A., Lankat-Buttgereit B., Göke B., Heimberg H., Nauck M.A., Schiemann U., Schatz H., Pfeiffer A.F.H.: Cloning and quantitative determination of the human Ca^{2+}/calmodulin-dependent protein kinase II *(CaMK II)* isoforms in human beta cells 465

Roll U., Turck C.W., Gitelman S.E., Rosenthal S.M., Nolte M.S., Masharani U., Ziegler A.-G., Baekkeskov S.: Peptide mapping and characterisation of glycation patterns of the glima 38 antigen recognised by autoantibodies in Type I diabetic patients 598

Rooman I., Heremans Y., Heimberg H., Bouwens L.: Modulation of rat pancreatic acinoductal transdifferentiation and expression of PDX-1 in vitro 907

Rowley K.G., Iser D.M., Best J.D., O'Dea K., Leonard D., McDermott R.: Albuminuria in Australian Aboriginal people: prevalence and associations with components of the metabolic syndrome 1397

Sackmann H., Tran-Van T., Tack I., Hanaire-Broutin H., Tauber J.-P., Ader J.-L.: Contrasting renal functional reserve in very long-term Type I diabetic patients with and without nephropathy 227

Saeki K., Xie J., Notkins A.L.: Genomic structure of mouse *IA-2*: comparison with its human homologue 1429

Sainsbury A., Wilks D., Cooney G.J.: Interaction between adrenal glucocorticoids and parasympathetic activation in mediating hyperinsulinaemia during long-term central neuropeptide Y infusion in rats 859

Sanai T., Sobka T., Johnson T., El-Essawy M., Muchaneta-Kubara E.C., Gharbia O.B., Oldroyd S., El Nahas A.M.: Expression of cytoskeletal proteins during the course of experimental diabetic nephropathy 91

Schrauwen P., Schaart G., Saris W.H.M., Slieker L.J., Glatz J.F.C., Vidal H., Blaak E.E.: The effect of weight reduction on skeletal muscle UCP2 and UCP3 mRNA expression and UCP3 protein content in Type II diabetic subjects 1408

Schütt M., Meier M., Meyer M., Klein J., Aries S.P., Klein H.H.: The HIV-1 protease inhibitor indinavir impairs insulin signalling in HepG2 hepatoma cells 1145

Shapira M.Y., Appelbaum E.Y., Hirshberg B., Mizrahi Y., Bar-On H., Ziv E.: A sustained, non-insulin related, hypoglycaemic effect of electroacupuncture in diabetic *Psammomys obesus* 809

Shin S.J., Lai F.J., Wen J.D., Hsiao P.J., Hsieh M.C., Tzeng T.F., Chen H.C., Guh J.Y., Tsai J.H.: Neuronal and endothelial nitric oxide synthase expression in outer medulla of streptozotocin-induced diabetic rat kidney 649

Sima A.A.F., Zhang W., Xu G., Sugimoto K., Guberski D., Yorek M.A.: A comparison of diabetic polyneuropathy in Type II diabetic BBZDR/Wor rats and in Type I diabetic BB/Wor rats 786

Sobel D.O., Ahvazi B., Jun H.-S., Chung Y.-H., Yoon J.-W.: Cyclophosphamide inhibits the development of diabetes in the diabetes-prone *BB* rat 986

Soedamah-Muthu S. S., Colhoun H. M., Taskinen M.-R., Idzior-Walus B., Fuller J. H. and the EURODIAB IDDM Complications Study Group: Differences in HDL-cholesterol: apoA-I + apoA-II ratio and apoE phenotype with albuminuric status in Type I diabetic patients 1353
Söhnlein P., Müller M., Syren K., Hartmann U., Böhm B. O., Meinck H. M., Knip M., Akerblom H. K., The Childhood Diabetes in Finland Study Group, Richter W.: Epitope spreading and a varying but not disease-specific GAD65 antibody response in Type I diabetes 210
Solini A., Chiozzi P., Falzoni S., Morelli A., Fellin R., Virgilio F. D.: High glucose modulates P2X7 receptor-mediated function in human primary fibroblasts 1248
Spranger J., Hammes H.-P., Preissner K. T., Schatz H., Pfeiffer A. F. H.: Release of the angiogenesis inhibitor angiostatin in patients with proliferative diabetic retinopathy: association with retinal photocoagulation 1404
Stefan N., Fritsche A., Madaus A., Häring H., Stumvoll M.: Stimulatory effect of increased non-esterified fatty acid concentrations on proinsulin processing in healthy humans 1368
Steinbrenner H., Lohmann T., Ostendorf B., Scherbaum W. A., Seissler J.: Autoantibodies to ICA12 (SOX-13) are not specific for Type I diabetes 1381
Stene L. C., Ulriksen J., Magnus P., Joner G.: Use of cod liver oil during pregnancy associated with lower risk of Type I diabetes in the offspring 1093
Strack V., Hennige A. M., Krützfeldt J., Bossenmaier B., Klein H.-H., Kellerer M., Lammers R., Häring H.-U.: Serine residues 994 and 1023/25 are important for insulin receptor kinase inhibition by protein kinase C isoforms β 2 and θ 443
Suarez-Pinzon W. L., Power R. F., Rabinovitch A.: Fas ligand-mediated mechanisms are involved in autoimmune destruction of islet beta cells in non-obese diabetic mice 1149
Suhonen L., Hiilesmaa V., Teramo K.: Glycaemic control during early pregnancy and fetal malformations in women with Type I diabetes mellitus 79
Suzuki E., Kashiwagi A., Hidaka H., Maegawa H., Nishio Y., Kojima H., Haneda M., Yasuda H., Morikawa S., Inubushi T., Kikkawa R.: ^{1}H- and ^{31}P-magnetic resonance spectroscopy and imaging as a new diagnostic tool to evaluate neuropathic foot ulcers in Type II diabetic patients 165
Suzuki E., Kashiwagi A., Nishio Y., Kojima H., Maegawa H., Haneda M., Yasuda H., Morikawa S., Inubushi T., Kikkawa R.: Usefulness of waveform analysis of popliteal artery in Type II diabetic patients using gated magnetic resonance 2D-cine-PC imaging and ^{31}P spectroscopy 1031
't Hart L. M., Dekker J. M., van Haeften T. W., Ruige J. B., Stehouwer C. D. A., Erkelens D. W., Heine R. J., Maassen J. A.: Reduced second phase insulin secretion in carriers of a sulphonylurea receptor gene variant associating with Type II diabetes mellitus 515
Tang J., Zhu X. W., Lust W. D., Kern T. S.: Retina accumulates more glucose than does the embryologically similar cerebral cortex in diabetic rats 1417
Tarnow L., Kjeld T., Knudsen E., Major-Pedersen A., Parving H.-H.: Lack of synergism between long-term poor glycaemic control and three gene polymorphisms of the renin angiotensin system on risk of developing diabetic nephropathy in Type I diabetic patients 794
The EURODIAB Substudy 2 Study Group: Infections and vaccinations as risk factors for childhood Type I (insulin-dependent) diabetes mellitus: a multicentre case-control investigation 47
Toutouzas K., Stefanadis C., Tsiamis E., Vlachopoulos Ch., Tousoulis D., Tsioufis C., Toutouzas P.: Aortic pressure-diameter relation in patients with non-insulin dependent diabetes mellitus: new insights 1070
Tree T. I. M., Morgenthaler N. G., Duhindan N., Hicks K. E., Madec A.-M., Scherbaum W. A., Banga J. P.: Two amino acids in glutamic acid decarboxylase act in concert for maintainance of conformational determinants recognised by Type I diabetic autoantibodies 881
Tripathy D., Carlsson Å.-L., Lehto M., Isomaa B., Tuomi T., Groop L.: Insulin secretion and insulin sensitivity in diabetic subgroups: studies in the prediabetic and diabetic state 1476
Ueda H., Ikegami H., Kawaguchi Y., Fujisawa T., Nojima K., Babaya N., Yamada K., Shibata M., Yamato E., Ogihara T.: Age-dependent changes in phenotypes and candidate gene analysis in a polygenic animal model of Type II diabetes mellitus; NSY mouse 932
Ueno N., Inui A., Asakawa A., Takao F., Tani S., Komatsu Y., Itoh Z., Kasuga M.: Erythromycin improves glycaemic control in patients with Type II diabetes mellitus 411
Vallejo S., Angulo J., Peiró C., Nevado J., Sánchez-Ferrer A., Petidier R., Sánchez-Ferrer C. F., Rodríguez-Mañas L.: Highly glycated oxyhaemoglobin impairs nitric oxide relaxations in human mesenteric microvessels 83
van Popele N. M., Westendorp I. C. D., Bots M. L., Reneman R. S., Hoeks A. P. G., Hofman A., Grobbee D. E., Witteman J. C. M.: Variables of the insulin resistance syndrome are associated with reduced arterial distensibility in healthy non-diabetic middle-aged women 665
Vasir B., Reitz P., Xu G., Sharma A., Bonner-Weir S., Weir G. C.: Effects of diabetes and hypoxia on gene markers of angiogenesis (HGF, cMET, uPA and uPAR, TGF-α, TGF-β, bFGF and Vimentin) in cultured and transplanted rat islets 763
Vauhkonen I., Niskanen L., Knip M., Ilonen J., Vanninen E., Kainulainen S., Uusitupa M., Laakso M.: Impaired insulin secretion in non-diabetic offspring of probands with latent autoimmune diabetes mellitus in adults 69
Vay D., Vidali M., Allochis G., Cusaro C., Rolla R., Mottaran E., Bellomo G., Albano E.: Antibodies against advanced glycation end product N^{ε}-(carboxymethyl)lysine in healthy controls and diabetic patients 1385
Velho G., Hattersley A. T., Froguel P.: Maternal diabetes alters birth weight in glucokinase-deficient (MODY2) kindred but has no influence on adult weight, height, insulin secretion or insulin sensitivity 1060
Villa M. P., Multari G., Montesano M., Pagani J., Cervoni M., Midulla F., Cerone E., Ronchetti R.: Sleep apnoea in children with diabetes mellitus: effect of glycaemic control 696
Weyer C., Foley J. E., Bogardus C., Tataranni P. A., Pratley R. E.: Enlarged subcutaneous abdominal adipocyte size, but not obesity itself, predicts Type II diabetes independent of insulin resistance 1498
Weyer C., Hanson K., Bogardus C., Pratley R. E.: Long-term changes in insulin action and insulin secretion associated with gain, loss, regain and maintenance of body weight 36
Yamada S., Zhu Q., Aihara Y., Onda H., Zhang Z., Yu L., Jin L., Si Y.-J., Nishigori H., Tomura H., Inoue I., Morikawa A., Yamagata K., Hanafusa T., Matsuzawa Y., Takeda J.: Cloning of cDNA and the gene encoding human hepatocyte nuclear factor (HNF)-3β and mutation screening in Japanese subjects with maturity-onset diabetes of the young 121
Yamaoka T., Yano M., Yamada T., Matsushita T., Moritani M., Ii S., Yoshimoto K., Hata J., Itakura M.: Diabetes and pancreatic tumours in transgenic mice expressing Pa × 6 332
Yang Q., Yamagata K., Yamamoto K., Cao Y., Miyagawa J., Fukamizu A., Hanafusa T., Matsuzawa Y.: R127W-HNF-4α is a loss of function mutation but not a rare polymorphism and causes Type II diabetes in a Japanese family with MODY1 520

Yaworsky K., Somwar R., Ramlal T., Tritschler H. J., Klip A.: Engagement of the insulin-sensitive pathway in the stimulation of glucose transport by α-lipoic acid in 3T3-L1 adipocytes 294
Yuan X., Yamada K., Ishiyama-Shigemoto S., Koyama W., Nonaka K.: Identification of polymorphic loci in the promoter region of the serotonin 5-HT2C receptor gene and their association with obesity and Type II diabetes 373
Yudkin J. S., Panahloo A., Stehouwer C., Emeis J. J., Bulmer K., Mohamed-Ali V., Denver A. E.: The influence of improved glycaemic control with insulin and sulphonylureas on acute phase and endothelial markers in Type II Diabetic subjects 1099
Zanetti M., Sato J., Katusic Z. S., O'Brien T.: Gene transfer of endothelial nitric oxide synthase alters endothelium-dependent relaxations in aortas from diabetic rabbits 340
Zhu Q., Yamagata K., Yu L., Tomura H., Yamada S., Yang Q., Yoshiuchi I., Sumi S., Miyagawa J., Takeda J., Hanafusa T., Matsuzawa Y.: Identification of missense mutations in the hepatocyte nuclear factor-3β gene in Japanese subjects with late-onset Type II diabetes mellitus 1197

For debate

Bergman R. N.: Non-esterified fatty acids and the liver: why is insulin secreted into the portal vein? 946
Richter B., Berger M.: Randomized controlled trials remain fundamental to clinical decision making in Type II diabetes mellitus: a comment to the debate on randomized controlled trials 254
Vijan S., Kent D. M., Hayward R. A.: Are randomized controlled trials sufficient evidence to guide clinical practice in Type II (non-insulin-dependent) diabetes mellitus? 125
Wiernsperger N.: Defects in microvascular haemodynamics during prediabetes: contributor or epiphenomenon? 1439

Letters to the editor

Atkinson M., Honeyman M., Peakman M., Roep B.: T-Cell markers in Type I diabetes: progress, prospects and realistic expectations 820
Bjørkhaug L., Søvik O., Bell G. I., Njølstad P. R., Molven A.: A simple test for the high-frequency P291fsinsC mutation in the HNF1α/MODY3 gene 818
Brownrigg L. M., Stanton K. G., McCann V. J., Burke V., Taylor R. R., van Bockxmeer F. M.: Increased frequency of the lipoprotein lipase 9N allele in adults with Type I (insulin-dependent) diabetes mellitus 953
Bruno G., Arcari R., Pagano A., Cerutti F., Berrino M., Pagano G.: Genetic heterogeneity by age at onset of Type I diabetes: higher prevalence of patients with 0 susceptible heterodimers in adults than in children in the registry of Turin, Italy 260
Bruno G., Biggeri A., Merletti F., Pagano G.: Temporal tends in incidence rates of Type I diabetes in Germany: birth cohort and calendar period effects 1334
Bulman M. P., Ellard S., Hattersley A. T.: R127Wis a loss-of-function mutation in HNF4α causing maturity-onset diabetes of the young (MODY) in a UK Caucasian family 1203
Chajara A., Raoudi M., Delpech B., Leroy M., Basuyau J. P., Levesque H.: Circulating hyaluronan and hyaluronidase are increased in diabetic rats 387
Cseh K., Winkler G., Melczer Z., Baranyi E.: The role of tumour necrosis factor (TNF)-α resistance in obesity and insulin resistance 525
Das U. N.: Possible beneficial action(s) of glucose-insulin-potassium regimen in acute myocardial infarction and inflammatory conditions: a hypothesis 1081
Dosch H.-M., Becker D. J.: Measurement of T-cell autoreactivity in autoimmune diabetes 386
Feltbower R. G., McKinney P. A., Bodansky H. J.: Rising incidence of childhood diabetes is seen at all ages and in urban and rural settings in Yorkshire, United Kingdom 682
Galama J. M. D.: Vaccinations as risk factors for Type I diabetes mellitus 684
Ilic S., Jovanovic L., Wollitzer A. O.: Is the paradoxical first trimester drop in insulin requirement due to an increase in C-peptide concentration in pregnant Type I diabetic women? 1329
Iwatani M., Wasada T., Katsumori K., Watanabe-Takahashi C., Kamatani N., Iwamoto Y.: Troglitazone decreases serum uric acid concentrations in Type II diabetic patients and non-diabetics 814
Jermendy G., Nádas J., Sápi Z.: "Lipoblastoma-like" lipotrophy induced by human insulin: morphological evidence for local dedifferentiation of adipocytes? 954
Kanauchi M., Nishioka H., Dohi K.: Secreted protein acidic and rich in cysteine (SPARC) in patients with diabetic nephropathy and tubulointerstitial injury 1076
Konrad D., Sobetzko D., Schmitt B., Schoenle E. J.: Insulin-dependent diabetes mellitus induced by the antitussive agent dextromethorphan 261
Kujala U. M., Kaprio J., Koskenvuo M.: Diabetes in a population-based series of twin pairs discordant for leisure sedentariness 259
Lefrandt J. D., Diercks G. F. H., van Boven A. J., Crijns H. J. G. M., van Gilst W. H., Gans R. O. B.: High fasting glucose and QTc duration in a large healthy cohort 1332
Lorini R., Avanzini M. A., Lenta E., Cotellessa M., De Giacomo C., d'Annunzio G.: Antibodies to tissue transglutaminase C in newly diagnosed and long-standing Type I diabetes mellitus 815
Lynn S., Hattersley A. T., McCarthy M. I., Frayling T. M., Turnbull D. M., Walker M.: Intermediate expansions of a X25/frataxin gene GAA repeat and Type II diabetes: assessment using parent-offspring trios 384
Mason S., Ong K. K. L., Pembrey M. E., The ALSPAC Study Team, Woods K. A., Dunger D. B.: The Gly972Arg variant in insulin receptor substrate-1 is not associated with birth weight in contemporary English children 1201
Matz R.: Sulphonylureas, magnesium and cardiac K^+ ATP channels 1080
Ng M. C. Y., Li J. K. Y., So W. Y., Critchley J. A. J. H., Cockram C. S., Bell G. I., Chan J. C. N.: Nature or nurture: an insightful illustration from a Chinese family with hepatocyte nuclear factor-1α diabetes (MODY3) 816
Pardini V. C., Pardini H., Velho G.: Accuracy of fasting glucose to diagnose diabetes in Brazilian subjects 132
Patterson C. C., Dahlquist G., Soltész G. (on behalf of the EURODIAB Substudy 2 Study Group) 684
Porte D. Jr., Seeley R. J., Woods S. C., Baskin D. G., Figlewicz D. P., Schwartz M. W.: Comments on the relation among GAD-65, IA-2 antibodies and body adiposity 133
Pozzilli P., Visalli N., Leslie D. and the IMDIAB Group: No evidence of rapid onset (Japanese) Type I diabetes in Caucasian patients 1332
Prisco F., Iafusco D., Franzese A., Sulli N., Barbetti F.: MODY 2 presenting as neonatal hyperglycaemia: a need to reshape the definition of "neonatal diabetes"? 1331
Ricart W., Bach C., Fernández-Real J. M., Sabrià J.: Major fetal complications in optimised pregestational diabetes mellitus 1077
Rique S., Ibáñez L., Marcos M. V., Carrascosa A., Potau N.: Effects of metformin on androgens and insulin concentrations in type A insulin resistance syndrome 385
Rosenbauer J., Herzig P., von Kries R., Giani G.: Temporal trends in incidence

rates of Type I diabetes in Germany: calendar period rather than birth-cohort effects 1335
Squires P. E., Persaud S. J., Jones P. M., Buchan A. M. J.: The extracellular calcium-sensing receptor in PHHI beta cells: does reduced autoinhibitory input contribute to hypersecretion of insulin? 1078
van Tilburg J. H. O., Rozeman L. B., van Someren H., Rigters-Aris C. A. E., Freriks J. P., Pearson P. L., Sandkuijl L. A., van Haeften T. W., Wijmenga C.: The exon 16–3*t* variant of the sulphonylurea receptor gene is not a risk factor for Type II diabetes mellitus in the Dutch Breda cohort 681
Walhor T., Schober E., Karimian-Teherani D., Rami B., and the Austrian Diabetes Incidence Study Group: Regional differences and temporal incidence trend of Type I diabetes mellitus in Austria from 1989 to 1999: a nationwide study 1449
Weng J. P., Lehto M., Forsblom C., Huang X., Li H., Groop L. C.: Hepatocyte nuclear factor-1β (MODY5) gene mutations in Scandinavian families with early-onset diabetes or kidney disease or both 131
Wookey P., Cooper M.: Amylin and hypertension: the status of the current debate 262

Book reviews

388, 1204

EASD News section

1, 17, 19, 25, 33, 35, 47, 53

Errata

526, 686, 1451

List of referees

1573

Author index

1586

Subject index

1585

Contents of Volume 43

I